


		240mm								
	120mm	120mm								
<p><b>HIGHLIGHTS OF PRESCRIBING INFORMATION</b></p> <p>These highlights do not include all the information needed to use NEOSTIGMINE METHYLSULFATE INJECTION safely and effectively. See full prescribing information for NEOSTIGMINE METHYLSULFATE INJECTION.</p> <p><b>NEOSTIGMINE METHYLSULFATE INJECTION, for intravenous use</b> Initial U.S. Approval: 1939</p> <p style="text-align: center;">-----INDICATIONS AND USAGE-----</p> <p>Neostigmine Methylsulfate Injection, a cholinesterase inhibitor, is indicated for the reversal of the effects of non-depolarizing neuromuscular blocking agents (NMBAs) after surgery (1).</p> <p style="text-align: center;">-----DOSAGE AND ADMINISTRATION-----</p> <ul style="list-style-type: none"> <li>Should be administered by trained healthcare providers (2.1)</li> <li>Peripheral nerve stimulator and monitoring for twitch responses should be used to determine when neostigmine methylsulfate injection should be initiated and if additional doses are needed (2.2) <ul style="list-style-type: none"> <li>For reversal of NMBAs with shorter half-lives, when first twitch response is substantially greater than 10% of baseline, or when a second twitch is present: <b>0.03 mg/kg by intravenous route</b> (2.2)</li> <li>For reversal of NMBAs with longer half-lives or when first twitch response is close to 10% of baseline: <b>0.07 mg/kg by intravenous route</b> (2.2)</li> </ul> </li> <li>Maximum total dosage is 0.07 mg/kg or up to a total of 5 mg (whichever is less) (2.2)</li> <li>An anticholinergic agent, e.g., atropine sulfate or glycopyrrolate, should be administered prior to or concomitantly with neostigmine methylsulfate injection (2.4)</li> </ul> <p style="text-align: center;">-----DOSAGE FORMS AND STRENGTHS-----</p> <p>Injection: 0.5 mg/mL and 1 mg/mL in 10 mL multiple-dose vials (3)</p> <p style="text-align: center;">-----CONTRAINDICATIONS-----</p> <ul style="list-style-type: none"> <li>Hypersensitivity to neostigmine (4)</li> <li>Peritonitis or mechanical obstruction of the intestinal or urinary tract (4)</li> </ul> <p style="text-align: center;">-----WARNINGS AND PRECAUTIONS-----</p> <ul style="list-style-type: none"> <li>Bradycardia: Atropine or glycopyrrolate should be administered prior to neostigmine methylsulfate to lessen risk of bradycardia.(5.1)</li> <li>Serious Reactions with Coexisting Conditions: Use with caution in patients with, coronary artery disease, cardiac arrhythmias, recent acute coronary syndrome or myasthenia gravis. (5.2)</li> <li>Neuromuscular Dysfunction: Can occur if large doses of neostigmine methylsulfate are administered when neuromuscular blockade is minimal; reduce dose if recovery from neuromuscular blockade is nearly complete. (5.4)</li> </ul> <p style="text-align: center;">-----ADVERSE REACTIONS-----</p> <p>Most common adverse reactions during treatment: bradycardia, nausea and vomiting. (6)</p> <p><b>To report SUSPECTED ADVERSE REACTIONS, contact Somerset Therapeutics, LLC at +1 800-417-9175 or FDA at 1-800-FDA-1088 or <a href="http://www.fda.gov/medwatch">www.fda.gov/medwatch</a></b></p> <p style="text-align: center;">-----USE IN SPECIFIC POPULATIONS-----</p> <ul style="list-style-type: none"> <li>Pregnancy: No human data and limited animal data exist. Use only if clearly needed.</li> </ul>	<p style="text-align: right;">Revised: 07/2021</p> <p><b>11. DESCRIPTION</b></p> <p>Neostigmine methylsulfate, a cholinesterase inhibitor, is (<i>m</i>-hydroxyphenyl) trimethylammonium methylsulfate dimethylcarbamate. The structural formula is:</p>  <p>Each mL of the 0.5 mg/mL strength contains neostigmine methylsulfate 0.5 mg, phenol 4.5 mg (used as preservative) and sodium acetate trihydrate 0.2 mg, in water for injection. The pH is adjusted, when necessary, with acetic acid/sodium hydroxide to achieve a value of 5.5.</p> <p>Each mL of the 1 mg/mL strength contains neostigmine methylsulfate 1 mg, phenol 4.5 mg (used as preservative), and sodium acetate trihydrate 0.2 mg, in water for injection. The pH is adjusted, when necessary, with acetic acid/sodium hydroxide to achieve a value of 5.5.</p> <p><b>12. CLINICAL PHARMACOLOGY</b></p> <p><b>12.1. Mechanism of Action</b> Neostigmine methylsulfate is a competitive cholinesterase inhibitor. By reducing the breakdown of acetylcholine, neostigmine methylsulfate induces an increase in acetylcholine in the synaptic cleft which competes for the same binding site as nondepolarizing neuromuscular blocking agents, and reverses the neuromuscular blockade.</p> <p><b>12.2. Pharmacodynamics</b> Neostigmine methylsulfate-induced increases in acetylcholine levels results in the potentiation of both muscarinic and nicotinic cholinergic activity. The resulting elevation of acetylcholine competes with nondepolarizing neuromuscular blocking agents to reverse neuromuscular blockade. Neostigmine methylsulfate does not readily cross the blood-brain barrier and, therefore, does not significantly affect cholinergic function in the central nervous system.</p> <p><b>12.3. Pharmacokinetics</b> <i>Distribution:</i> Following intravenous injection, the observed neostigmine methylsulfate volume of distribution is reported between 0.12 and 1.4 L/kg. Protein binding of neostigmine methylsulfate to human serum albumin ranges from 15 to 25 %.</p> <p><i>Metabolism:</i> Neostigmine methylsulfate is metabolized by microsomal enzymes in the liver.</p> <p><i>Elimination:</i> Following intravenous injection, the reported elimination half-life of neostigmine methylsulfate is between 24 and 113 minutes. Total body clearance of neostigmine methylsulfate is reported between 1.14 and 16.7 mL/min/kg.</p> <p><i>Renal impairment:</i> Elimination half-life of neostigmine methylsulfate was prolonged in anephric patients compared to normal subjects; elimination half-life for normal, transplant and anephric patients were 79.8 ± 48.6, 104.7 ± 64 and 181 ± 54 min (mean ± SD), respectively.</p> <p><i>Hepatic impairment:</i> The pharmacokinetics of neostigmine methylsulfate in patients with hepatic impairment has not been studied.</p> <p><i>Pediatrics:</i> Elimination half-life of neostigmine methylsulfate in infants (2-10 months), children (1-6 years) and adults (29-48 years) were 39 ± 5 min, 48 ± 16 min, and 67 ± 8 min (mean ± SD), respectively. Observed neostigmine methylsulfate clearance for infants, children and adults were 14 ± 3, 11 ± 3 and 10 ± 2 mL/min/kg (mean ± SD), respectively.</p> <p><i>Drug Interaction Studies:</i> The pharmacokinetic interaction between neostigmine methylsulfate and other drugs has not been studied.</p> <p><b>13. NONCLINICAL TOXICOLOGY</b></p> <p><b>13.1. Carcinogenesis, Mutagenesis, Impairment of Fertility</b> <i>Carcinogenesis:</i> Long-term animal studies have not been performed to evaluate the carcinogenic potential of neostigmine.</p> <p><i>Genotoxicity:</i> Neostigmine methylsulfate was not mutagenic or clastogenic when evaluated in an in vitro bacterial reverse mutation assay (Ames test), an in vitro Chinese hamster ovary cell chromosomal aberration assay, or an in vivo mouse bone marrow micronucleus assay.</p> <p><i>Impairment of Fertility:</i> In a fertility and early embryonic development study in rats, male rats were treated for 28 days prior to mating and female rats were treated for 14 days prior to mating with intravenous neostigmine methylsulfate (human equivalent doses of 1.6, 4, and 8.1 mcg/kg/day, based on body surface area). No adverse effects were reported at any dose (up to 0.1- times the MRHD of 5 mg/60 kg person based on a body surface area comparison).</p> <p><b>14. CLINICAL STUDIES</b></p> <p>The evidence for the efficacy of neostigmine methylsulfate for the reversal of the effects of non-depolarizing neuromuscular blocking agents after surgery is derived from the published literature. Randomized, spontaneous-recovery or placebo-controlled studies using similar efficacy endpoints evaluated a total of 404 adult and 80 pediatric patients undergoing various surgical procedures. Patients had reductions in their recovery time from neuromuscular blockade with neostigmine methylsulfate treatment compared to spontaneous recovery.</p> <p><b>16. HOW SUPPLIED/STORAGE AND HANDLING</b></p> <p>Neostigmine methylsulfate injection, USP is a clear, colorless liquid available in the following presentations:</p> <table border="1"> <thead> <tr> <th>NDC No.</th> <th>Strength</th> <th>Vial size</th> </tr> </thead> <tbody> <tr> <td>70069-805-10</td> <td>0.5 mg/mL</td> <td>10 mL multiple-dose vials (supplied in packages of 10)</td> </tr> <tr> <td>70069-806-10</td> <td>1 mg/mL</td> <td>10 mL multiple-dose vials (supplied in packages of 10)</td> </tr> </tbody> </table> <p>The vial stopper is not made with natural rubber latex.</p> <p>Neostigmine methylsulfate injection, USP should be stored at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light. Store in carton until time of use.</p> <p><b>Manufactured for:</b> Somerset Therapeutics, LLC Hollywood, FL 33024 <b>Customer Care # 1-800-417-9175</b></p> <p><b>Manufactured by:</b> Caplin Steriles Limited, Gummidiipoondi - 601 201,</p> <p>Made in India Code: TN/Drugs/TN00003457</p> <p style="text-align: right;">Revised: 07/2021 13160939</p>	NDC No.	Strength	Vial size	70069-805-10	0.5 mg/mL	10 mL multiple-dose vials (supplied in packages of 10)	70069-806-10	1 mg/mL	10 mL multiple-dose vials (supplied in packages of 10)
NDC No.	Strength	Vial size								
70069-805-10	0.5 mg/mL	10 mL multiple-dose vials (supplied in packages of 10)								
70069-806-10	1 mg/mL	10 mL multiple-dose vials (supplied in packages of 10)								
<p><b>FULL PRESCRIBING INFORMATION: CONTENTS*</b></p> <p><b>1. INDICATIONS AND USAGE</b></p> <p><b>2. DOSAGE AND ADMINISTRATION</b></p> <p>2.1. Important Dosage Information</p> <p>2.2. Dosage in Adults</p> <p>2.3. Dosage in Pediatric Patients, including Neonates</p> <p>2.4. Anticholinergic (Atropine or Glycopyrrolate) Administration</p> <p><b>3. DOSAGE FORMS AND STRENGTHS</b></p> <p><b>4. CONTRAINDICATIONS</b></p> <p><b>5. WARNINGS AND PRECAUTIONS</b></p> <p>5.1. Bradycardia</p> <p>5.2. Serious Adverse Reactions in Patients with Certain Coexisting Conditions</p> <p>5.3. Hypersensitivity</p> <p>5.4. Neuromuscular Dysfunction</p> <p>5.5. Cholinergic Crisis</p> <p><b>6. ADVERSE REACTIONS</b></p> <p>6.1. Clinical Trials Experience</p> <p>6.2. Post Marketing Experience</p> <p><b>7. DRUG INTERACTIONS</b></p> <p><b>FULL PRESCRIBING INFORMATION</b></p> <p><b>1. INDICATIONS AND USAGE</b></p> <p>Neostigmine methylsulfate injection is a cholinesterase inhibitor indicated for the reversal of the effects of non-depolarizing neuromuscular blocking agents after surgery.</p> <p><b>2. DOSAGE AND ADMINISTRATION</b></p> <p><b>2.1. Important Dosage Information</b> Neostigmine methylsulfate injection should be administered by trained healthcare providers familiar with the use, actions, characteristics, and complications of neuromuscular blocking agents (NMBA) and neuromuscular block reversal agents. Doses of neostigmine methylsulfate injection should be individualized, and a peripheral nerve stimulator should be used to determine the time of initiation of neostigmine methylsulfate injection and should be used to determine the need for additional doses.</p> <p>Neostigmine methylsulfate injection is for intravenous use only and should be injected slowly over a period of at least 1 minute. The neostigmine methylsulfate injection dosage is weight-based [see <i>Dosage and Administration</i> (2.2)].</p> <p>Prior to neostigmine methylsulfate injection administration and until complete recovery of normal ventilation, the patient should be well ventilated and a patent airway maintained. Satisfactory recovery should be judged by adequacy of skeletal muscle tone and respiratory measurements in addition to the response to peripheral nerve stimulation.</p> <p>An anticholinergic agent, e.g., atropine sulfate or glycopyrrolate, should be administered prior to or concomitantly with neostigmine methylsulfate injection [see <i>Dosage and Administration</i> (2.4)].</p> <p><b>2.2. Dosage in Adults</b></p> <ol style="list-style-type: none"> <li>Peripheral nerve stimulation devices capable of delivering a train-of-four (TOF) stimulus are essential to effectively using neostigmine methylsulfate injection.</li> <li>There must be a twitch response to the first stimulus in the TOF of at least 10% of its baseline level, i.e., the response prior to NMBA administration, prior to the administration of neostigmine methylsulfate injection.</li> <li>Prior to administration, visually inspect neostigmine methylsulfate injection for particulate matter and discoloration.</li> <li>Neostigmine methylsulfate injection should be injected slowly by intravenous route over a period of at least 1 minute.</li> <li>A 0.03 mg/kg to 0.07 mg/kg dose of neostigmine methylsulfate will generally achieve a TOF twitch ratio of 90% (TOF<sub>90%</sub>) within 10 to 20 minutes of administration. Dose selection should be based on the extent of spontaneous recovery that has occurred at the time of administration, the half-life of the NMBA being reversed, and whether there is a need to rapidly reverse the NMBA.</li> </ol> <ul style="list-style-type: none"> <li>The 0.03 mg/kg dose is recommended for: <ul style="list-style-type: none"> <li>Reversal of NMBAs with shorter half-lives, e.g., rocuronium, or</li> <li>When the first twitch response to the TOF stimulus is substantially greater than 10% of baseline or when a second twitch is present.</li> </ul> </li> <li>The 0.07 mg/kg dose is recommended for: <ul style="list-style-type: none"> <li>NMBAs with longer half-lives, e.g., vecuronium and pancuronium, or</li> <li>When the first twitch response is relatively weak, i.e., not substantially greater than 10% of baseline or</li> <li>There is need for more rapid recovery.</li> </ul> </li> </ul> <ol style="list-style-type: none"> <li>TOF monitoring should continue to be used to evaluate the extent of recovery of neuromuscular function and the possible need for an additional dose of neostigmine methylsulfate injection.</li> <li>TOF monitoring alone should not be relied upon to determine the adequacy of reversal of neuromuscular blockade as related to a patient's ability to adequately ventilate and maintain a patent airway following tracheal extubation.</li> <li>Patients should continue to be monitored for adequacy of reversal from NMBAs for a period of time that would assure full recovery based on the patient's medical condition and the pharmacokinetics of neostigmine and the NMBA used.</li> <li>The recommended maximum total dose is 0.07 mg/kg or up to a total of 5 mg, whichever is less.</li> </ol> <p><b>2.3. Dosage in Pediatric Patients, including Neonates</b> Adult guidelines should be followed when neostigmine methylsulfate injection is administered to pediatric patients. Pediatric patients require neostigmine methylsulfate injection doses similar to those for adult patients.</p> <p><b>2.4. Anticholinergic (Atropine or Glycopyrrolate) Administration</b> An anticholinergic agent, e.g., atropine sulfate or glycopyrrolate, should be administered prior to or concomitantly with neostigmine methylsulfate injection. The anticholinergic agent should be administered intravenously using a separate syringe. In the presence of bradycardia, it is recommended that the anticholinergic agent be administered prior to neostigmine methylsulfate injection.</p> <p><b>3. DOSAGE FORMS AND STRENGTHS</b></p> <p>Neostigmine methylsulfate injection is available as</p> <ul style="list-style-type: none"> <li>Injection: 0.5 mg/mL, 5 mg of neostigmine methylsulfate in 10 mL multiple-dose vials</li> <li>Injection: 1 mg/mL, 10 mg of neostigmine methylsulfate in 10 mL multiple-dose vials</li> </ul> <p><b>8. USE IN SPECIFIC POPULATIONS</b></p> <p><b>8.1. Pregnancy</b> Risk Summary There are no adequate or well-controlled studies of neostigmine methylsulfate in pregnant women. It is not known whether neostigmine methylsulfate can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. The incidence of malformations in human pregnancies has not been established for neostigmine as the data are limited. All pregnancies, regardless of drug exposure, have a background risk of 2 to 4 % for major birth defects, and 15 to 20 % for pregnancy loss.</p> <p>No adverse effects were noted in rats or rabbits treated with human equivalent doses of neostigmine methylsulfate doses up to 8.1 and 13 mcg/kg/day, respectively, during organogenesis (0.1 to 0.2-times the maximum recommended human dose of 5 mg/60 kg person/day based on body surface area comparisons).</p> <p>Anticholinesterase drugs, including neostigmine may cause uterine irritability and induce premature labor when administered to pregnant women near term.</p> <p>Neostigmine methylsulfate should be given to a pregnant woman only if clearly needed.</p> <p><b>Data</b> <i>Animal Data</i> In embryofetal development studies, rats and rabbits were administered neostigmine methylsulfate at human equivalent doses (HED, on a mg/m<sup>2</sup> basis) of 1.6, 4 and 8.1 mcg/kg/day and 3.2, 8.1, and 13 mcg/kg/day, respectively, during the period of organogenesis (Gestation Days 6 through 17 for rats and Gestation Days 6 through 18 for rabbits). There was no evidence for a teratogenic effect in rats and rabbits up to HED 8.1 and 13 mcg/kg/day, which are approximately 0.097 times and 0.16 times the MRHD of 5 mg/60 kg, respectively in the presence of minimal maternal toxicity (tremors, ataxia, and prostration). The studies resulted in exposures in the animals well below predicted exposures in humans.</p> <p>In a pre- and postnatal development study in rats, neostigmine methylsulfate was administered to pregnant female rats at human equivalent doses (HED) of 1.6, 4 and 8.1 mcg/kg/day from Day 6 of gestation through Day 20 of lactation, with weaning on Day 21. There were no adverse effects on physical development, behavior, learning ability, or fertility in the offspring at HED doses up to 8.1 mcg/kg/day which is 0.097 times the MRHD of 5 mg/60 kg on a mg/m<sup>2</sup> basis in the presence of minimal maternal toxicity (tremors, ataxia, and prostration). The studies resulted in exposures in the animals well below predicted exposures in humans.</p> <p><b>8.2. Lactation</b> <i>Risk Summary.</i> Neostigmine methylsulfate has not been studied in lactating women. It is not known whether neostigmine methylsulfate is present in human milk, or if neostigmine methylsulfate has effects on milk production or the breastfed child. Therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's need for neostigmine methylsulfate and any potential adverse effects on the breastfed child from neostigmine methylsulfate or from the underlying maternal condition.</p> <p><b>8.4. Pediatric Use</b> Neostigmine methylsulfate is approved for the reversal of the effects of non-depolarizing neuromuscular blocking agents after surgery in pediatric patients of all ages.</p> <p>Recovery of neuromuscular activity occurs more rapidly with smaller doses of cholinesterase inhibitors in infants and children than in adults. However, infants and small children may be at greater risk of complications from incomplete reversal of neuromuscular blockade due to decreased respiratory reserve. The risks associated with incomplete reversal outweigh any risk from giving higher doses of neostigmine methylsulfate (up to 0.07 mg/kg or up to a total of 5 mg, whichever is less).</p> <p>The dose of neostigmine methylsulfate required to reverse neuromuscular blockade in children varies between 0.03 mg - 0.07 mg/kg, the same dose range shown to be effective in adults, and should be selected using the same criteria as used for adult patients [see <i>Clinical Pharmacology</i> (12.3)].</p> <p>Since the blood pressure in pediatric patients, particularly infants and neonates, is sensitive to changes in heart rate, the effects of an anticholinergic agent (e.g., atropine) should be observed prior to administration of neostigmine to lessen the probability of bradycardia and hypotension.</p> <p><b>8.5. Geriatric Use</b> Because elderly patients are more likely to have decreased renal function, neostigmine methylsulfate should be used with caution and monitored for a longer period in elderly patients. The duration of action of neostigmine methylsulfate is prolonged in the elderly; however, elderly patients also experience slower spontaneous recovery from neuromuscular blocking agents. Therefore, dosage adjustments are not generally needed in geriatric patients; however, they should be monitored for longer periods than younger adults to assure additional doses of neostigmine methylsulfate are not required. The duration of monitoring should be predicated on the anticipated duration of action for the NMBA used on the patient [see <i>Dosage and Administration</i> (2.3)].</p> <p><b>8.6. Renal Impairment</b> Elimination half-life of neostigmine methylsulfate was prolonged in anephric patients compared to normal subjects.</p> <p>Although no adjustments to neostigmine methylsulfate dosing appear to be warranted in patients with impaired renal function, they should be closely monitored to assure the effects of the neuromuscular blocking agent, particularly one cleared by the kidneys, do not persist beyond those of neostigmine methylsulfate. In this regard, the interval for re-dosing the neuromuscular blocking agent during the surgical procedure may be useful in determining whether, and to what extent, post-operative monitoring needs to be extended.</p> <p><b>8.7. Hepatic Impairment</b> The pharmacokinetics of neostigmine methylsulfate in patients with hepatic impairment have not been studied. Neostigmine methylsulfate is metabolized by microsomal enzymes in the liver. No adjustments to the dosing of neostigmine methylsulfate appear to be warranted in patients with hepatic insufficiency. However, patients should be carefully monitored if hepatically cleared neuromuscular blocking agents were used during their surgical procedure as their duration of action may be prolonged by hepatic insufficiency whereas neostigmine methylsulfate, which undergoes renal elimination, will not likely be affected. This could result in the effects of the neuromuscular blocking agent outlasting those of neostigmine methylsulfate. This same situation may arise if the neuromuscular blocking agent has active metabolites. In this regard, the interval for re-dosing the neuromuscular blocking agent during the surgical procedure may be useful in determining whether, and to what extent, post-operative monitoring needs to be extended.</p> <p><b>10. OVERDOSAGE</b></p> <p>Muscarinic symptoms (nausea, vomiting, diarrhea, sweating, increased bronchial and salivary secretions, and bradycardia) may appear with overdose of neostigmine methylsulfate (cholinergic crisis), but may be managed by the use of additional atropine or glycopyrrolate. The possibility of iatrogenic overdose can be lessened by carefully monitoring the muscle twitch response to peripheral nerve stimulation. Should overdose occur, ventilation should be supported by artificial means until the adequacy of spontaneous respiration is assured, and cardiac function should be monitored.</p> <p>Overdosage of neostigmine methylsulfate can cause cholinergic crisis, which is characterized by increasing muscle weakness, and through involvement of the muscles of respiration, may result in death. Myasthenic crisis, due to an increase</p>										