




Pemetrexed 500 mg Carton - ANDA

Size: 52 mm x 52 mm x 86 mm (LxWxH)

Colours:

- Pantone Process Blue C
- Pantone 300 C
- Black

<div>■ Pantone 300 C</div> <div>■ Black</div>											
<div>500 mg/vial</div> <div>Pemetrexed for Injection, USP</div> <div>Discard Unused Portion</div> <div>Single-Dose Vial</div> <div>NDC 70069-835-01</div>			<div>1200030350</div> <div>V00</div>								
			<div>NDC 70069-835-01</div>								
			<div>Code No.: MH/DRUGS/28-KD/476</div>								
<div>Pemetrexed for Injection, USP</div>			<div>Pemetrexed for Injection, USP</div>								
<div>Pemetrexed for Injection, USP</div>			<div>Pemetrexed for Injection, USP</div>								
<div>Usual Dosage: See accompanying literature for dosage.</div> <div>Storage: Store powder at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].</div> <div>Store reconstituted and infusion solutions at refrigerated, 2-8°C (36-46°F).</div> <div>Caution: Cytotoxic Agent</div> <div></div> <div>3 70069 83501 2</div>			<div>Each vial contains pemetrexed disodium equivalent to 500 mg pemetrexed, 500 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.</div> <div>To Reconstitute: Add 20 mL of 0.9% Sodium Chloride Injection, USP (preservative free) to make a solution containing 25 mg/mL pemetrexed. Reconstituted solution must be further diluted before use (see accompanying literature).</div> <div>Administer infusion solution within 24 hours after initial reconstitution. Discard unused portion.</div>			<div>500 mg/vial</div> <div>For intravenous use only.</div> <div>Rx only</div> <div>Single-Dose Vial</div> <div>Discard Unused Portion</div> <div></div>			<div>Serialization FPO</div> <div></div> <div>Manufactured for: Somerset Therapeutics, LLC. Somerset, NJ 08873</div> <div>Made in India</div>		